

Math 33 Solving a System of Three Equations in Three Unknowns Feb. 1, 2018

We will be solving systems of linear equations using a variety of techniques. To reduce the amount of writing in class, IU have provided worked out solutions to the exercises I will be completing in class. In this way, you can allow me (this time) to focus on the arithmetic while you focus on the process.

Consider the following system:

$$\begin{aligned}2x - 2y - 4z &= -6 \\2x + y - z &= -3 \\x + y + 2z &= 5\end{aligned}$$

I. The Elimination (or Math D) Method:

We will use the first two equations to eliminate the variable x and arrive at a single equation in only y and z . Multiplying the second equation by -1 and adding, we have:

$$\begin{array}{r}2x - 2y - 4z = -6 \\-2x - y + z = 3 \\ \hline-3y - 3z = -3\end{array}\quad (1)$$

We will now use the last two equations to eliminate the variable x and arrive at a second equation in only y and z . Multiplying the second equation by -2 and adding, we have:

$$\begin{array}{r}2x + y - z = -3 \\-2x - 2y - 4z = -10 \\ \hline-y - 5z = -13\end{array}\quad (2)$$

Using the equations (1) and (2), we will now consider an easier system of two equations in two unknowns:

$$\begin{aligned}-3y - 3z &= -3 \\-y - 5z &= -13\end{aligned}$$

Multiplying both sides of the first equation by $-\frac{1}{3}$, then adding, we can eliminate the variable y from the system and arrive at an equation solely in z .

$$\begin{array}{r}y + z = 1 \\-y - 5z = -13 \\ \hline-4z = -12 \\z = 3\end{array}\quad (3.1)$$

We will now find the value for the variable y by using our value for z and substituting back into equation (2):

$$\begin{aligned} -y - 5 \cdot 3 &= -13 \\ -y - 15 &= -13 \\ -y &= 2 \\ y &= -2 \quad (3.2) \end{aligned}$$

Finally, we will now use our determined values for both y and z , substitute them into one of the original three equations (we will use the third, for simplicity), and determine the value for the one remaining variable x :

$$\begin{aligned} x + (-2) + 2 \cdot 3 &= 5 \\ x + 4 &= 5 \\ x &= 1 \quad (3.3) \end{aligned}$$

From the equations (3.1), (3.2), and (3.3), we now have the ordered triple solution: $(1, -2, 3)$

II. Gauss-Jordan Elimination on an Augmented Matrix:

Rewriting our original system as an augmented matrix, we will apply a sequence of elementary row operations to rewrite that augmented matrix in reduced row-echelon form:

$$\begin{aligned} &\left[\begin{array}{ccc|c} 2 & -2 & -4 & -6 \\ 2 & 1 & -1 & -3 \\ 1 & 1 & 2 & 5 \end{array} \right] \\ \frac{1}{2}R_1 \rightarrow R_1 &\left[\begin{array}{ccc|c} 1 & -1 & -2 & -3 \\ 2 & 1 & -1 & -3 \\ 1 & 1 & 2 & 5 \end{array} \right] \\ -2R_1 + R_2 \rightarrow R_2 &\left[\begin{array}{ccc|c} 1 & -1 & -2 & -3 \\ 0 & 3 & 3 & 3 \\ -R_1 + R_3 \rightarrow R_3 & \left[\begin{array}{ccc|c} 1 & -1 & -2 & -3 \\ 0 & 3 & 3 & 3 \\ 0 & 2 & 4 & 8 \end{array} \right] \\ \frac{1}{3}R_2 \rightarrow R_2 &\left[\begin{array}{ccc|c} 1 & -1 & -2 & -3 \\ 0 & 1 & 1 & 1 \\ 0 & 2 & 4 & 8 \end{array} \right] \\ R_2 + R_1 \rightarrow R_1 &\left[\begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 1 & 1 \\ -2R_2 + R_3 \rightarrow R_3 & \left[\begin{array}{ccc|c} 1 & 0 & -1 & -2 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 2 & 6 \end{array} \right] \end{array} \right. \end{aligned}$$

$$\frac{1}{2}R_3 \rightarrow R_3 \begin{bmatrix} 1 & 0 & -1 & | & -2 \\ 0 & 1 & 1 & | & 1 \\ 0 & 0 & 1 & | & 3 \end{bmatrix}$$

$$\begin{array}{l} R_3 + R_1 \rightarrow R_1 \\ -R_3 + R_2 \rightarrow R_2 \end{array} \begin{bmatrix} 1 & 0 & 0 & | & 1 \\ 0 & 1 & 0 & | & -2 \\ 0 & 0 & 1 & | & 3 \end{bmatrix}$$

translating back from augmented matrix form to equation form, this gives us an equivalent system of three equations in the same three unknowns, specifically:

$$\begin{aligned} x &= 1 \\ y &= -2 \\ z &= 3 \end{aligned}$$

Again, we have a solution that is the ordered triple $(1, -2, 3)$

Now let us consider two additional techniques for solving a system of three equations in three unknowns:

III. Using a Single Matrix Equation:

We can rewrite the original system of three equations in three unknowns as a single matrix equation where A represents the matrix of coefficients of our three variables x , y , and z . The matrix \bar{x} , will be a 3×1 matrix (or equivalently a column vector) consisting of these three variables and \bar{b} will be a 3×1 matrix containing the constants that reside on the right-hand sides of the equations in the original system.

So our system can be generically rewritten in the form:

$$A\bar{x} = \bar{b}$$

Of course, our solution to this equation is:

$$\bar{x} = A^{-1}\bar{b}$$

With our system, we can rewrite as a single matrix equation:

$$\begin{bmatrix} 2 & -2 & -4 \\ 2 & 1 & -1 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -6 \\ -3 \\ 5 \end{bmatrix}$$

whose solution is:

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 & -2 & -4 \\ 2 & 1 & -1 \\ 1 & 1 & 2 \end{bmatrix}^{-1} \begin{bmatrix} -6 \\ -3 \\ 5 \end{bmatrix} \quad (*)$$

Unfortunately, we cannot multiply on the right-hand side of this equation without first taking a break and finding A^{-1} .

Aside:

$$\begin{aligned} & \begin{bmatrix} 2 & -2 & -4 & | & 1 & 0 & 0 \\ 2 & 1 & -1 & | & 0 & 1 & 0 \\ 1 & 1 & 2 & | & 0 & 0 & 1 \end{bmatrix} \\ R_1 \leftrightarrow R_3 & \begin{bmatrix} 1 & 1 & 2 & | & 0 & 0 & 1 \\ 2 & 1 & -1 & | & 0 & 1 & 0 \\ 2 & -2 & -4 & | & 1 & 0 & 0 \end{bmatrix} \\ -2R_1 + R_2 \rightarrow R_2 & \begin{bmatrix} 1 & 1 & 2 & | & 0 & 0 & 1 \\ 0 & -1 & -5 & | & 0 & 1 & -2 \\ -2R_1 + R_3 \rightarrow R_3 & \begin{bmatrix} 1 & 1 & 2 & | & 0 & 0 & 1 \\ 0 & -1 & -5 & | & 0 & 1 & -2 \\ 0 & -4 & -8 & | & 1 & 0 & -2 \end{bmatrix} \\ -R_2 \rightarrow R_2 & \begin{bmatrix} 1 & 1 & 2 & | & 0 & 0 & 1 \\ 0 & 1 & 5 & | & 0 & -1 & 2 \\ 0 & -4 & -8 & | & 1 & 0 & -2 \end{bmatrix} \\ -R_2 + R_1 \rightarrow R_1 & \begin{bmatrix} 1 & 0 & -3 & | & 0 & 1 & -1 \\ 0 & 1 & 5 & | & 0 & -1 & 2 \\ 4R_2 + R_3 \rightarrow R_3 & \begin{bmatrix} 1 & 0 & -3 & | & 0 & 1 & -1 \\ 0 & 1 & 5 & | & 0 & -1 & 2 \\ 0 & 0 & 12 & | & 1 & -4 & 6 \end{bmatrix} \\ \frac{1}{12}R_3 \rightarrow R_3 & \begin{bmatrix} 1 & 0 & -3 & | & 0 & 1 & -1 \\ 0 & 1 & 5 & | & 0 & -1 & 2 \\ 0 & 0 & 1 & | & \frac{1}{12} & -\frac{1}{3} & \frac{1}{2} \end{bmatrix} \\ 3R_3 + R_1 \rightarrow R_1 & \begin{bmatrix} 1 & 0 & 0 & | & \frac{1}{4} & 0 & \frac{1}{2} \\ 0 & 1 & 0 & | & -\frac{5}{12} & \frac{2}{3} & -\frac{1}{2} \\ -5R_3 + R_2 \rightarrow R_2 & \begin{bmatrix} 1 & 0 & 0 & | & \frac{1}{4} & 0 & \frac{1}{2} \\ 0 & 1 & 0 & | & -\frac{5}{12} & \frac{2}{3} & -\frac{1}{2} \\ 0 & 0 & 1 & | & \frac{1}{12} & -\frac{1}{3} & \frac{1}{2} \end{bmatrix} \end{aligned}$$

Now that we have found that $A^{-1} = \begin{bmatrix} \frac{1}{4} & 0 & \frac{1}{2} \\ -\frac{5}{12} & \frac{2}{3} & -\frac{1}{2} \\ \frac{1}{12} & -\frac{1}{3} & \frac{1}{2} \end{bmatrix}$, we can multiply on the

right-hand side of the equation (*):

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 & -2 & -4 \\ 2 & 1 & -1 \\ 1 & 1 & 2 \end{bmatrix}^{-1} \begin{bmatrix} -6 \\ -3 \\ 5 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1/4 & 0 & 1/2 \\ -5/12 & 2/3 & -1/2 \\ 1/12 & -1/3 & 1/2 \end{bmatrix} \begin{bmatrix} -6 \\ -3 \\ 5 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \cdot (-6) + 0 \cdot (-3) + \frac{1}{2} \cdot 5 \\ -\frac{5}{12} \cdot (-6) + \frac{2}{3} \cdot (-3) - \frac{1}{2} \cdot 5 \\ \frac{1}{12} \cdot (-6) - \frac{1}{3} \cdot (-3) + \frac{1}{2} \cdot 5 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}$$

Yet again, we have a solution that is the ordered triple $(1, -2, 3)$

For our final attempt at solving this system of three equations in three unknowns, we will employ what is referred to as *Cramer's Rule*, which employs the use of determinants:

IV. Cramer's Rule:

Let us first compute D , the determinant of the coefficient matrix for the system of equations:

$$\begin{aligned} D &= \begin{vmatrix} 2 & -2 & -4 \\ 2 & 1 & -1 \\ 1 & 1 & 2 \end{vmatrix} = 2 \cdot \begin{vmatrix} 1 & -1 \\ 1 & 2 \end{vmatrix} - (-2) \cdot \begin{vmatrix} 2 & -1 \\ 1 & 2 \end{vmatrix} + (-4) \cdot \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} \\ &= 2 \cdot 3 - (-2) \cdot 5 + (-4) \cdot 1 \\ &= 6 + 10 - 4 \\ &= 12 \end{aligned}$$

Replacing the first column (the column consisting of the coefficients for the variable x in each of our three equations) of the determinant D with the constants on the right-hand side of the equal signs in each of the original equations of our system, we can compute D_x :

$$\begin{aligned}
 D_x &= \begin{vmatrix} -6 & -2 & -4 \\ -3 & 1 & -1 \\ 5 & 1 & 2 \end{vmatrix} = -6 \cdot \begin{vmatrix} 1 & -1 \\ 1 & 2 \end{vmatrix} - (-2) \cdot \begin{vmatrix} -3 & -1 \\ 5 & 2 \end{vmatrix} + (-4) \cdot \begin{vmatrix} -3 & 1 \\ 5 & 1 \end{vmatrix} \\
 &= -6 \cdot 3 - (-2) \cdot (-1) + (-4) \cdot (-8) \\
 &= -18 - 2 + 32 \\
 &= 12
 \end{aligned}$$

Repeating this process on the second and third columns, we arrive at:

$$\begin{aligned}
 D_y &= \begin{vmatrix} 2 & -6 & -4 \\ 2 & -3 & -1 \\ 1 & 5 & 2 \end{vmatrix} = 2 \cdot \begin{vmatrix} -3 & -1 \\ 5 & 2 \end{vmatrix} - (-6) \cdot \begin{vmatrix} 2 & -1 \\ 1 & 2 \end{vmatrix} + (-4) \cdot \begin{vmatrix} 2 & -3 \\ 1 & 5 \end{vmatrix} \\
 &= 2 \cdot (-1) - (-6) \cdot 5 + (-4) \cdot 13 \\
 &= -2 + 30 - 52 \\
 &= -24
 \end{aligned}$$

and

$$\begin{aligned}
 D_z &= \begin{vmatrix} 2 & -2 & -6 \\ 2 & 1 & -3 \\ 1 & 1 & 5 \end{vmatrix} = 2 \cdot \begin{vmatrix} 1 & -3 \\ 1 & 5 \end{vmatrix} - (-2) \cdot \begin{vmatrix} 2 & -3 \\ 1 & 5 \end{vmatrix} + (-6) \cdot \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} \\
 &= 2 \cdot 8 - (-2) \cdot 13 + (-6) \cdot 1 \\
 &= 16 + 26 - 6 \\
 &= 36
 \end{aligned}$$

Finally, we have that $x = \frac{D_x}{D} = \frac{12}{12} = 1$,

$$y = \frac{D_y}{D} = \frac{-24}{12} = -2, \text{ and}$$

$$z = \frac{D_z}{D} = \frac{36}{12} = 3$$

And one last time, we arrive at the solution that is the ordered triple $(1, -2, 3)$